



Draft Genome Sequence of *Vibrio parahaemolyticus* VH3, Isolated from an Aquaculture Environment in Greece

Castillo, Daniel; Jun, Jin Woo; D'Alvise, Paul; Middelboe, Mathias; Gram, Lone; Liu, Siyang; Katharios, Pantelis

Published in:
Genome Announcements

Link to article, DOI:
[10.1128/genomeA.00731-15](https://doi.org/10.1128/genomeA.00731-15)

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Castillo, D., Jun, J. W., D'Alvise, P., Middelboe, M., Gram, L., Liu, S., & Katharios, P. (2015). Draft Genome Sequence of *Vibrio parahaemolyticus* VH3, Isolated from an Aquaculture Environment in Greece. *Genome Announcements*, 3(4), [e00731-15]. <https://doi.org/10.1128/genomeA.00731-15>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Draft Genome Sequence of *Vibrio parahaemolyticus* VH3, Isolated from an Aquaculture Environment in Greece

Daniel Castillo,^a Jin Woo Jun,^b Paul D'Alvise,^c Mathias Middelboe,^a Lone Gram,^c Siyang Liu,^d Pantelis Katharios^b

Marine Biological Section, University of Copenhagen, Helsingør, Denmark^a; Institute of Marine Biology, Biotechnology and Aquaculture, Hellenic Centre for Marine Research, Crete, Greece^b; Department of Systems Biology, Technical University of Denmark, Copenhagen, Denmark^c; BGI Europe A/S, Copenhagen, Denmark^d

D.C. and J.W.J. contributed equally to this work.

***Vibrio parahaemolyticus* is an important foodborne pathogen responsible for gastroenteritis outbreaks globally. It has also been identified as an important pathogen in aquatic organisms. Here, we report a draft genome sequence of *V. parahaemolyticus*, strain VH3, isolated from farmed juvenile greater amberjack, *Seriola dumerili*, in Greece.**

Received 27 May 2015 Accepted 28 May 2015 Published 2 July 2015

Citation Castillo D, Jun JW, D'Alvise P, Middelboe M, Gram L, Liu S, Katharios P. 2015. Draft genome sequence of *Vibrio parahaemolyticus* VH3, isolated from an aquaculture environment in Greece. *Genome Announc* 3(4):e00731-15. doi:10.1128/genomeA.00731-15.

Copyright © 2015 Castillo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Pantelis Katharios, katharios@hcmr.gr.

Vibrio parahaemolyticus is one of the most important foodborne pathogens globally. Infections are almost exclusively associated with the consumption of raw or improperly cooked contaminated seafood (1, 2). Clinical characteristics include watery diarrhea, abdominal cramps, nausea, and fever (3, 4). However, *V. parahaemolyticus* infections are not exclusive to humans and have recently been demonstrated to be a major source of disease in aquaculture production (5, 6). Therefore, monitoring of this pathogenic *Vibrio* species in aquaculture environments is important for both the aquaculture industry and human health. In the present study, *V. parahaemolyticus* VH3 was isolated from the aquaculture facility of the Institute of Aquaculture of the Hellenic Centre for Marine Research (HCMR) at the port of Heraklion, Greece, during a vibriosis outbreak in juvenile greater amberjack.

Genomic DNA of *V. parahaemolyticus* VH3 was extracted using the QIAamp DNA minikit (Qiagen) according to the manufacturer's protocol. The genome was sequenced by standard shotgun sequencing methods using a 454 GS-FLX Titanium sequencing system (Roche). The sequence data consisted of 5,600,000 reads (average length, 79 bp), providing 89-fold coverage. *De novo* assembly of the whole sequencing reads was performed with a GS De Novo Assembler. A total of 67 contigs, with a minimum length of 943 bp and a maximum length of 592,284 bp (N_{50} , 187.7 kbp), were obtained. Annotation was performed by the NCBI Prokaryotic Genome Automatic Annotation Pipeline (PGAAP) (7). Additionally, the genome was screened for presence of specific genetic markers for *V. parahaemolyticus* (*tlh*, *tdh*, *trh*, *orf8*, and *toxRSnew*) (8), prophages (9), virulence factors (10), and antibiotic resistance genes (11).

The draft genome of *V. parahaemolyticus* VH3 was 4,955,051 bp in length, with a G+C composition of 45.3%. Genome annotation resulted in 4,338 coding sequences (CDS), 39 tRNAs, 37 pseudogenes, and 4 rRNAs. One phage-related sequence of 12.5 kb was found. Virulence factors related to adhesion (arylsulfatases and colonization factor Acfa), hemo-

lysin, metalloproteases, and *Mycobacterium* virulence operon and multidrug resistance efflux pumps were found. Antibiotic resistant genes were detected for fluoroquinolones and tetracycline.

To the best of our knowledge, this is the first genome report of a *V. parahaemolyticus* strain isolated in an aquaculture system in Greece. The current sequence data generated here will contribute to the understanding of genome variability of *V. parahaemolyticus* isolates in future genomic studies.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at GenBank under the accession number [LCVL000000000](https://www.ncbi.nlm.nih.gov/nuclink/LCVL000000000). The version described in this paper is version LCVL01000000.

ACKNOWLEDGMENTS

This work was supported by a grant from the Danish Directorate for Food, Fisheries and Agri Business (ProAqua, project 09-072829), by FP7, Marie Curie IRSES 2010 (AQUAPHAGE, project 269175), and by FISHPHAGE project 131, funded by the European Social Fund and Greek National resources under the "Excellence" scheme, NSRF 2007-2013.

REFERENCES

- Daniels NA, MacKinnon L, Bishop R, Altekruse S, Ray B, Hammond RM, Thompson S, Wilson S, Bean NH, Griffin PM, Slutsker L. 2000. *Vibrio parahaemolyticus* infections in the United States, 1973–1998. *J Infect Dis* 181:1661–1666. <http://dx.doi.org/10.1086/315459>.
- Jun JW, Kim HJ, Yun SK, Chai JY, Park SC. 2014. Eating oysters without risk of vibriosis: application of a bacteriophage against *Vibrio parahaemolyticus* in oysters. *Int J Food Microbiol* 188:31–35. <http://dx.doi.org/10.1016/j.ijfoodmicro.2014.07.007>.
- Matsumoto C, Okuda J, Ishibashi M, Iwanaga M, Garg P, Rammamurthy T, Wong HC, Depaula A, Kim YB, Albert MJ, Nishibuchi M. 2000. Pandemic spread of an O3:K6 clone of *Vibrio parahaemolyticus* and emergence of related strains evidenced by arbitrarily primed PCR and *toxRS* sequence analyses. *J Clin Microbiol* 38:578–585.
- Ottaviani D, Leoni F, Rocchegiani E, Santarelli S, Canonico C, Masini L, Ditrani V, Carraturo A. 2008. First clinical report of pandemic *Vibrio parahaemolyticus* O3:K6 infection in Italy. *J Clin Microbiol* 46:2144–2145. <http://dx.doi.org/10.1128/JCM.00683-08>.

5. Soto-Rodriguez SA, Gomez-Gil B, Lozano-Olvera R, Betancourt-Lozano M, Morales-Covarrubias MS. 2015. Field and experimental evidence of *Vibrio parahaemolyticus* as the causative agent of acute hepatopancreatic necrosis disease of cultured shrimp (*Litopenaeus vannamei*) in northwestern Mexico. *Appl Environ Microbiol* 81:1689–1699. <http://dx.doi.org/10.1128/AEM.03610-14>.
6. Tey YH, Jong KJ, Fen SY, Wong HC. 2015. Occurrence of *Vibrio parahaemolyticus*, *Vibrio cholerae*, and *Vibrio vulnificus* in the aquacultural environments of Taiwan. *J Food Prot* 78:969–976. <http://dx.doi.org/10.4315/0362-028X.JFP-14-405>.
7. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciufo S, Li W. 2013. Prokaryotic genome annotation pipeline. National Center for Biotechnology Information, Bethesda, MD.
8. García K, Torres R, Uribe P, Hernández C, Rioseco ML, Romero J, Espejo RT. 2009. Dynamics of clinical and environmental *Vibrio parahaemolyticus* strains during seafood-related summer diarrhea outbreaks in southern Chile. *Appl Environ Microbiol* 75:7482–7487. <http://dx.doi.org/10.1128/AEM.01662-09>.
9. Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool. *Nucleic Acids Res* 39:W347–W352. <http://dx.doi.org/10.1093/nar/gkr485>.
10. Joensen KG, Scheut F, Lund O, Hasman H, Kaas RS, Nielsen EM, Aarestrup FM. 2014. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 52:1501–1510. <http://dx.doi.org/10.1128/JCM.03617-13>.
11. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <http://dx.doi.org/10.1093/jac/dks261>.